

Joint media response concerning “the discovery of a new kind of killer whale” off western North America

For immediate release

October 8th, 2021. 3:00 PM Pacific Time

On October 4th, Hakai Magazine published an article with a headline reporting the discovery of “a new kind of killer whale”. Other media outlets subsequently reported similar headlines in Canada, the USA and internationally, and although some content has since been corrected in these articles, others continue to report false information.

The fact is that no new type of killer whale has been discovered and that many of the whales in question are part of the known Bigg’s (transient) killer whale population that prey on marine mammals. Their distribution patterns, social behaviours and population structure have been studied for over 30 years by several government and non-government research organizations in California, Washington, British Columbia, and Alaska.

Some previous publications on these killer whales include a 1997 catalogue by Black et al.¹ which provides details on 105 individuals photographed off the outer coast of California. It presents information on 45 of the 150 whales documented in the recent study by McInnes et al.² referenced by the media. Most of these whales have been photo-identified nearly each year since 1997, many have reproduced, and the overall population size has grown to around 200 whales (Black et al. in prep). A book about killer whales published in 1999 by Ford and Ellis³ states that some of these same whales had also been observed in British Columbia. Then in 2013, Ford et al.⁴ went on to designate some 217 of these killer whales as belonging to a putative “outer-coast” sub-population and 304 belonging to a putative inner-coast sub-population. In 2019, Towers et al.⁵ updated these numbers to 241 and 381, respectively, but confirmed that there is immigration and emigration between the two putative sub-populations.

Published records of a more oceanic population of mammal-eating killer whales off the west coast of North America also exist. Black et al.¹ reported on killer whales photographed far offshore of California that had cookiecutter shark scars and stated that these whales “may belong to currently unknown types”. Hoelzel et al.⁶ presented genetic evidence in 1998 “indicating that there is another population found off California” and that “their genotype was closest to the transient type”. Pitman et al.⁷ suggested in 2001 that a group of mammal-eating killer whales documented in the open ocean off California may “belong to a far offshore population that normally occurs seaward of the continental shelf”. Finally, Towers⁸ communicated in 2017 that he and colleagues had encountered Bigg’s killer whales far offshore of British Columbia “that looked and behaved differently from any others we had known”. These four publications independently suggest that more than one population of Bigg’s killer whales occur off the west coast of North America. The new study by McInnes et al.² also states that some of the whales in their catalogue “may belong to an undescribed pelagic offshore mammal-eating population” but do not go on to discuss or conclude the existence of any population beyond the previously described and putative outer-coast population.

The terminology used to describe these killer whale populations can sometimes be confusing. For example, the putative outer-coast population of Bigg's (transient) killer whales that eats mammals tends to travel along the shelf edge from Monterey Bay in California to Haida Gwaii in British Columbia, but occasionally comes into inner coastal waters throughout this range^{4,5}. A different population long referred to as offshore killer whales is only known to eat fish, mainly sharks⁹, and travels along the shelf edge from southern California to the Aleutian Islands¹⁰. They too will occasionally come into inshore waters. These populations are already both well described, do not mix, and are genetically distinct¹¹. Less is known about the so far unnamed population of mammal-eating killer whales that occur in the open ocean.

The large media response to this topic of interest highlights an appetite for advances in our understanding of killer whales globally. The recent study² quoted in these articles provides a valuable contribution of data on individual killer whales that are found in deep water off the west coast of North America; however, it does not provide new facts or interpretations about populations or their structure that are different to those already published by others.

Sincerely,

Jared Towers
Executive Director, Bay Cetology, info@baycetology.org



BAY CETOLOGY

Nancy Black
Director of Research, California Killer Whale Project, montereybaywhalewatch@gmail.com

and

Alisa Schulman-Janiger
Lead Research Biologist, California Killer Whale Project, asjaniger@gmail.com



Ken Balcomb
President, Center for Whale Research, kenbalcomb@gmail.com



Lance Barrett-Lennard
Director, Marine Mammal Research Program, Ocean Wise, Lance.barrett-lennard@ocean.org



Dan Olsen
Research Biologist, North Gulf Oceanic Society, waterdogdan@gmail.com



Howard Garrett
President, Orca Network, info@orcانetwork.org



Deborah Giles
Research Director, Wild Orca, giles@wildorca.org



John Ford
Scientist Emeritus, Fisheries and Oceans Canada, johnkbford@gmail.com



References

- 1: Black, N.A., Schulman-Janiger, A., Ternullo, R.L., and Guerrero-Ruiz, M. 1997. Killer Whales of California and Western Mexico: a Catalog of Photo-identified Individuals. NOAA-TM-NMFS-SWFSC-247.
- 2: McInnes, J.D., Mathieson, C.R., West-Stap, P.J., Marcos, S.L., Wade, V.L., Olson, P.A. and Trites, A.W. 2021. Transient killer whales of central and northern California and Oregon: A catalogue of photo-identified individuals. NOAA-TM-NMFS-SWFSC-644.
- 3: Ford, J.K.B., and G.M. Ellis. 1999. Transients: Mammal-Hunting Killer Whales. UBC Press, Vancouver, British Columbia. 96 pp.
- 4: Ford, J.K.B, Stredulinsky, E.H., Towers J.R. and Ellis, G.M. 2013. Information in Support of the Identification of Critical Habitat for Transient Killer Whales (*Orcinus orca*) off the West Coast of Canada. DFO Can. Sci. Advis. Sec. Res. Doc. 2012/155. iv + 46 p.
- 5: Towers, J.R., Sutton, G.J., Shaw, T.J.H., Malleson, M., Matkin, D., Gisborne, B., Forde, J., Ellifrit, D., Ellis, G.M., Ford, J.K.B., and Doniol-Valcroze, T. 2019. Photo-identification Catalogue, Population Status, and Distribution of Bigg's Killer Whales known from Coastal Waters of British Columbia, Canada. Can. Tech. Rep. Fish. Aquat. Sci. 3311: vi + 299 p.
- 6: Hoelzel, A.R., Dahlheim, M.E., and Stern, S.J. 1998. Low genetic variation among killer whales (*Orcinus orca*) in the eastern North Pacific and genetic differentiation between foraging specialists. J Hered. 89:121-128.
- 7: Pitman, R.L., Ballance, L.T., Mesnick, S.I., and Chivers, S.J. 2001. Killer whale predation on sperm whales: observations and implications. Mar. Mamm. Sci. 17:494-507.
- 8: Towers, J.R. 2017. Bigg's orcas: not so "transient" anymore. Orcazine.
- 9: Ford, J.K.B., Ellis, G.M., Matkin, C.O., Wetklo, M.H., Barrett-Lennard, L.G., and Withler, R.E. 2011. Shark predation and tooth wear in a population of northeastern Pacific killer whales. Aquat. Biol. 11:213-224.
- 10: Dahlheim, M.E., Schulman-Janiger, A., Black, N., Ternullo, R., Ellifrit, D., and Balcomb, K.C. 2008. Eastern temperate North Pacific offshore killer whales (*Orcinus orca*): occurrence, movements, and insights into feeding ecology. Mar. Mamm. Sci. 24:719–729.
- 11: Morin, P.A., Archer, F.I., Foote, A.D., Vilstrup, J., Allen, E.E., Wade, P., Durban, J., Parsons, K., Pitman, R., Li, L., Bouffard, P., Abel Nielsen, S.C., Rasmussen, M., Willerslev, E., Gilbert, M.T., and Harkins, T. 2010. Complete mitochondrial genome phylogeographic analysis of killer whales (*Orcinus orca*) indicates multiple species. Genome Research, 20:908– 916.